

High retention rate of Carlin tags and no effects on growth and condition of PIT tagging - a long-term field study on European eel

Elin Myrenäs^{*} , Joacim Näslund , Philip Jacobson , Birgitta Jacobson , John Persson ,
Håkan Wickström , Jennie Strömquist  and Josefin Sundin 

Department of Aquatic Resources, Institute of Freshwater Research, Swedish University of Agricultural Sciences, Stångholmsvägen 2, SE-17893, Drottningholm, Sweden

Received 1 December 2025 / Accepted 10 June 2026

Handling Editor: Nicolas BEZ

Abstract – Different types of tags and markers are commonly used for various fish monitoring and tracking purposes. Effects of tags and markers on fish and the retention rates can affect the interpretation of mark-recapture data on both the individual (e.g. growth and body condition) and population level (e.g. survival and production estimates), making studies of this issue important. In this study, we examined tag retention of Carlin tags and potential effects on growth and body condition of Passive Integrated Transponder (PIT)-tags on the European eel (*Anguilla anguilla*). We used data from several long-term tagging projects conducted in a natural Swedish lake, combining outlet trap monitoring with catch data from a commercial fisher. The longest of these studies spanned over 27 years, covering the majority of the eel's resident life stage. We found that the retention rate of Carlin tags was 92%, with 64% of the Carlin-tagged silver eels being recaptured. The recapture rate of PIT-tagged eels released as juveniles in the lake was 12%. This recapture rate indicates little or no effect on survival when compared to other studies of both tagged and untagged eels. No major impact of PIT-tags on growth and condition in eels was found. We conclude that both Carlin and PIT-tags are suitable methods for tagging European eel, each with strengths in its respective area of use.

Keywords: *Anguilla anguilla* / long-term monitoring / mark-recapture / Sweden / tag shedding

1 Introduction

Various tags and markers for individual identification have been important tools in scientific studies and management of fish for over a century, and the methods are constantly developing (Atkins, 1876; Latour, 2005; Lucas and Barras, 2008). Present-day physical tags and markers make it possible to study behaviour (Lucas and Barras, 2008), migration patterns (e.g. Acou et al., 2005; Piper et al., 2013; Wright et al., 2015), habitat use (e.g. Riley et al., 2011; Herrera et al., 2019), abundance (Cahill et al., 2018), and mortality (Vandergoot and Brenden, 2014) at the individual (identity marking) or group (batch marking) level. Tagging studies can also generate valuable information for stock assessments (Saville and Morrison, 1985; Höhne et al., 2023), and decision-making in fisheries and conservation management (Hays et al., 2019). At the same time, empirical evidence of the effects of tags and tagging procedures on fish remains limited and sometimes inconsistent. Some authors suggest that this is generally not a problem if fish are not too small (e.g. Nyqvist et al., 2023;

Závorka et al., 2024). Nevertheless, several studies have shown that tags and markers can cause adverse effects on fish, for example, in altering feeding patterns (Bridger and Booth, 2003), behaviour (Huusko et al., 2016), growth rate, and mortality (Larsen et al., 2013). Fish with conspicuous external tags might also experience higher predation pressure than untagged fish (Kerstetter et al., 2004). The handling and tagging procedure varies among tagging techniques, and some of the most invasive techniques are associated with a risk of inducing inflammation (Semple et al., 2018) and infection (Elliott and Pascho, 2001). In addition to the potential negative impacts of tagging on the studied fish, tag shedding can also affect the result of the study and lead to biased interpretations and inferences if not accounted for (Cowen and Schwarz, 2005). Currently, there is a smorgasbord of tagging methods available to enable individual identification of fish, which all come with their own set of pros and cons (Latour, 2005; Thorstad et al., 2013). Some methods suit certain species better than others; therefore, species-specific evaluations are needed to enable informed decisions on what method to use.

Tagging is particularly valuable for research and management of the European eel (*Anguilla anguilla*), due to their long life span and long-distance migration undertaken both as

^{*}Corresponding author: elin.myrenas@slu.se

larvae/juveniles and adults (Schmidt, 1912; Miller et al., 2019). It is with the use of varying tagging methods that scientists have been able to study the migrating mature eels on their way to spawn in the Sargasso Sea (Tesch, 1989; Westerberg et al., 2014; Wright et al., 2022). Since the European eel has a complex life cycle (Tesch, 1977) it is important to manage the species throughout all its life stages to support and secure its continued existence, given that the European eel is classified as Critically Endangered (Pike et al., 2020).

Two types of tags have been widely used in studies of yellow and silver eels: external Carlin tags (e.g., Westin, 1990; Pedersen and Dieperink, 2000; Sjöberg et al., 2009; Westerberg and Sjöberg, 2014), and internal Passive Integrated Transponder tags (PIT-tags) (e.g., Holmgren and Mosegaard, 1996; Laffaille et al., 2005; Riley et al., 2011; Wickström and Sjöberg, 2014). The Carlin tag (sometimes referred to as trailer tag, Eames and Hino, 1983) was developed in the 1950s and consists of a small plastic plate with an ID number and additional text information that is attached to the fish with two steel wires (Carlin, 1955). It is usually attached straight through the eel dorsal musculature. In Sweden, more than 3000 eels have been tagged with Carlin tags just in the past five years (data from SLU Aqua), and data on recaptured Carlin tagged silver eels reported by commercial fishers are used in the stock assessment to assess fishing mortality, “*F*” (Dekker and Sjöberg, 2013; van Gemert et al., 2024). Since the fishing mortality assessment is dependent on reported recaptures, it is influenced by tag shedding, should it occur. The retention rate of Carlin tags has been reported to be relatively low in field studies of some species, for example, Atlantic sturgeon (*Acipenser oxyrinchus*) and nurse shark (*Ginglymostoma cirratum*) (28% and 21%, respectively) (Carrier, 1985; Kallemeyn, 1989). In other species, the retention is high, for example, Chinook salmon (*Oncorhynchus tshawytscha*) and Atlantic salmon (*Salmo salar*) (98–99% and 98% respectively) (Eames and Hino, 1983; Huusko et al., 2016). Given these species-specific differences in retention rates of Carlin tags, assessing retention rates specifically for eels is pivotal for accurate evaluation of recapture data. Currently, such data is largely missing, but tag retention of Carlin tags in eel is considered to be high based on expert opinion (Nielsen, 1988). In addition to the expert opinion, we are aware of only one study, which is a laboratory experiment reporting a Carlin tag retention rate of ca 90% (Vøllestad, 1988).

The other main tag type used in eel, the passive integrated transponder (PIT-tag), was developed in the early 1980s (Prentice and Park, 1984). These tags have an electronic microchip with a unique ID code, enclosed in a glass hull (Prentice et al., 1990). The microchip remains inactive until it comes within range of a PIT-reader/antennae that powers the transponder, and the reader/antennae thereby receives the unique ID code via radio waves (Prentice et al., 1990). PIT-tags have a wide application field, ranging from individual identification of experimental fish (Holmgren and Mosegaard, 1996) and mark-recapture studies (Wickström and Sjöberg, 2014), to active movement biotelemetry in the field using portable PIT-antennas (Cucherousset et al., 2005) and passive movement and migration telemetry using larger stationary PIT-antennas (Riley et al., 2011). In eel, it is usually inserted in the abdominal cavity or the dorsal musculature (Tab. 1).

Tag retention of PIT-tags in eel is well studied and can be considered relatively high (Tab. 1), but there are still ambiguities to investigate regarding the effects of PIT-tagging (or implants of micro acoustic tags of comparable size, shape, and construction) on growth and mortality rates in eel. Although such effects have been specifically investigated in laboratories, mesocosms, and field studies, reported results range from no effects to major growth reductions in PIT-tagged eels (Tab. 1). Given the broad range of previous results, continued testing, for example, under differing environmental conditions, is important and a direct necessity to facilitate reliable meta-analyses on the effects of PIT-tags on European eels.

This study investigated the retention rate of Carlin tags and the potential effects on growth and condition of PIT-tagging in the European eel, using previously unpublished data from long-term tagging projects (both Carlin and PIT-tagging) in a natural Swedish lake. Unlike most previous studies on PIT-tagged eels, which are typically based on short-term experiments lasting weeks to months, the present work is based on data collected over an extended time span of up to 27 years. By combining long-term monitoring with a lake-based field setting, this study addresses a knowledge gap, as PIT-tagging research has largely been confined to controlled environments or riverine systems. We expected high tag retention for Carlin tagged silver eels in the field based on expert opinion and a previous laboratory study (Nielsen, 1988; Vøllestad, 1988). Based on the available literature, we did not expect PIT-tagging to negatively affect growth or condition (Tab. 1). All individuals, both tagged and control eels, originated from the same commercial catch.

2 Material and methods

Throughout the material and methods section, we report the author’s initials to clarify contributor roles for reproducibility and replicability using the Method Reporting with Initials for Transparency (MeRIT) guidelines (Nakagawa et al., 2023). In this paper, we used previously unpublished data from several projects related to monitoring of European eel in Sweden. This implies that the methods had a monitoring focus, rather than being designed as a specific study. Given the timespan of the data reported here, substantial efforts were made to locate and compile all necessary information and data (performed by JSu, BJ, PJ, EM, HW, JSt, and JP). The general background of the studies, methods, and data hosting has been published in a technical report (Jacobson et al., 2024).

2.1 Study site

The lake Ymsen (WGS84 dec: 58.666417, 13.965861) is a small and shallow lowland lake (average depth: 2.4 m, max depth: 4 m, total surface area: 13 km²) with little vegetation, mainly mud on the bottom, and some rocks and stones in the shore zone. Based on temperature measurements conducted four times annually (in February, April/May, August and October) between 1995 and 2025, water temperatures ranged from 0.1 to 23.1 °C, following seasonal air temperature changes (Miljödata-MVM, 2026). There are no significant tributaries, and the lake drains through one small river,

Table 1. Summary of results from studies on effects of Passive Integrated Transponder (PIT)-tagging on growth and survival, and PIT-tag retention in *Anguilla* species. Data on eel size at tagging is given as it was presented in the paper (i.e., if it says, “silver eel”, then no more specific information on size was provided in the paper). Placement of the tag is noted in the column “Retention” (AC – abdominal cavity; DM – Dorsal musculature (DMD – Dorsal musculature (DMD – dorsal fin; DMH – behind the head)). “NR” indicates missing methodological information (Not Reported); “NA” indicates parameters not investigated in the study.

References	Species	Location	Method	Tag length	Eel size	Study duration	Growth	Survival	Retention
Acou et al., 2000	<i>A. anguilla</i>	Mesocosm	NR	NR	Silver eel	30 days	NA	100%	100% (NR)
Morrison and Secor, 2003	<i>A. rostrata</i>	Field	Needle	NR	457 mm (mean)	2 months	NA	NA	89% (AC)
Verdon and Desrochers, 2003	<i>A. rostrata</i>	Field	Needle	14 mm	196–726 mm (mean ± SD: 472 ± 79 mm)	<16 months	NA	NA	98.6% (DMH)
Verdon et al., 2003	<i>A. rostrata</i>	Field	Needle	12–14 mm	380 mm (mean)	<3 months	NA	NA	93.9% (DMD)
Acou et al., 2005	<i>A. anguilla</i>	Lab	Needle	NR	>200 mm	1 hour	NA	NA	86% (AC)
Zimmerman and Welsh, 2008*	<i>A. rostrata</i>	Lab	Needle	12.5 mm	205–370 mm	9 weeks	NA	NA	100% (AC)
									100% (DMD)
									88% (DMH)
									>95% (AC)
Hirt-Chabbert and Young, 2012	<i>A. australis</i>	Lab	Scalpel	11 mm	101 ± 12 g (mean ± SD)	108 days	No effect	No effect	NA (AC)
Mazel et al., 2013	<i>A. anguilla</i>	Field	Trocar	12 mm	>200 mm	1–11 years	Reduced	NA	100% (NR)
Mazel et al., 2013	<i>A. anguilla</i>	Corf	Trocar	12 mm	247–732 mm	28 days	NA	NA	100% (AC)
Wright et al., 2015	<i>A. anguilla</i>	Stream enclosure	Scalpel	23 mm	356.0 ± 20.1 mm (mean ± SD)	7–14 days	NA	100%	100% (AC)
Jellyman and Crow, 2016	<i>A. australis</i>	Lab	Needle	15 mm	428–570 mm (mean: 646 mm)	21 days	NA	NA	100% (AC)
Jellyman and Crow, 2016	<i>A. australis</i>	Field	Needle	15 mm	>400 mm	7 days	NA	NA	99.1% (AC)
Jellyman and Crow, 2016	<i>A. australis</i>	Field	Needle	15 mm	>400 mm	9 months	NA	NA	95% (AC)
Schmucker et al., 2017	<i>A. rostrata</i>	Lab	Needle	12 mm	Silver eel	5 days	NA	No apparent mortality	100% (DMD)
Nzau Matondo and Ovidio, 2018	<i>A. anguilla</i>	Lab	Scalpel	23 mm	215–441 mm	20 days	NA	100%	100% (AC)
Herrera et al., 2019	<i>A. anguilla</i>	Lab	NR	8–12 mm	300 ± 10 mm (mean ± SD)	9 months	No effect	99%	100% (NR)
Nzau Matondo et al., 2021	<i>A. anguilla</i>	Field	Scalpel	12 mm	2–31.7 g (mean: 8.6 g)	Up to 2 years	Possibly reduced	NA	NA (AC)
Jepsen et al., 2022	<i>A. anguilla</i>	Ponds	Scalpel	12 mm	16.6–25.1 mm (mean: 21.6 mm)	76 days	No effect (non-significant)	97%; No effect	99% (AC)

* Alternatively, see Zimmerman (2008).

Table 2. Descriptive statistics of European eel (*Anguilla anguilla*) tagged with Carlin or Passive Integrated Transponder (PIT) tags and released in Lake Ymsen, Sweden. The table summarizes release date, tagging size (small yellow, large yellow eel, or silver eel), total length in range and mean (± 1 SD), total mass in range and mean (± 1 SD), number of eels tagged with PIT-tags, number of eels tagged with Carlin tags, and origin (caught in Lake Ymsen, or imported from England).

Date released	Tagging size	Length (mm), range; mean (± 1 SD)	Mass (g), range; mean (± 1 SD)	N PIT	N carlin	Origin
1998-10-28	Small yellow	171–297; 221 (± 23)	8–43; 17 (± 6)	190		England
1999-07-29	Large yellow	251–682; 542 (± 109)	21–521; 255 (± 130)	83		Lake Ymsen
1999-09-15	Small yellow	114–299; 207 (± 23)	5–41; 14 (± 5)	1512		England
2000-09-14	Small yellow	121–328; 231 (± 31)	3–71; 20 (± 10)	1087		England
2012-10-11	Silver eel	609–1030; 859 (± 68)	452–2120; 1276 (± 298)	118	118	Lake Ymsen

Ölebäcken (WGS84 dec: 58.645019, 13.934244). On the way towards the coast, the water from Ölebäcken flows to the river Tidan and onwards downstream via the much larger lake Vänern and the river Göta älv, where there are several migration barriers for eels (natural as well as human-made). The lake is therefore lacking open, natural fish migration routes, making it a well-controlled natural system suitable for eel monitoring. There are, however, eels in the lake due to translocation of naturally recruiting juvenile eels, caught in an eel trap/collector in the river Göta Älv (located next to an upstream migration barrier at Trollhättan, WGS84 dec: 58.274333, 12.271389), and due to restocking of imported juvenile eels from Europe (from Denmark, England, and France). This translocation and restocking of juvenile eel in Ymsen have been ongoing for at least a century (Nyström and Trybom, 1902; Nordqvist, 1928; van Gemert et al., 2024), with notes on eel presence dating back to at least 1883 (Vallin, 1929).

2.2 PIT- and Carlin tagging

Between 1998 and 2012, five separate tagging events were conducted in the lake Ymsen, with various sizes of eels and with two kinds of tags, either with only PIT- tags (tagging event 1–4), or with both PIT- tags and Carlin tags (tagging event 5).

During tagging events 1–3, small yellow eels were PIT-tagged each fall (September–October), between the years 1998 and 2000, and released in Lake Ymsen (1998: $N=190$, mean size = 221 mm; 1999: $N=1527$, mean size = 207 mm; 2000: $N=1100$, mean size = 231 mm; Tab. 2). The eels had been purchased from a commercial supplier of imported quarantined glass eels for restocking and aquaculture purposes (Scandinavian Silver Eel AB (SSE), Helsingborg, Sweden; eel origin: Severn, England). Prior to release, the imported eels were held in quarantine at an aquaculture facility from spring to fall each year to ensure they had reached a sufficient size to minimize adverse effects from PIT-tagging (Jepsen et al., 2005). The eels were tagged with PIT-tags (12 mm, 0.1 g, Trovan Ltd, Great Britain) at the SSE facilities (by HW and other staff at the Swedish Board of Fisheries). The PIT-tagging was performed under sedation with 0.12 g L⁻¹ Benzocaine (C₉H₁₁NO₂, Sigma-Aldrich, Inc., St. Louis, Missouri, USA; dissolved in a small amount of 95% ethanol before being mixed

in the bath of water). The PIT-tag was inserted using a syringe/needle or by hand into the abdominal cavity after making a small incision at the side of the ventral midline, a few centimetres anterior to the anus, using a scalpel blade. This method was chosen because scalpel incision is easier and safer than using needle injectors alone to penetrate the skin for tagging of eels due to their soft abdomen and tough skin, which increases the risk of internal organ damage when using needles (Zimmerman and Welsh, 2008; Hirt-Chabbert and Young, 2012). Data on total length (± 0.5 mm), total mass (± 0.05 g) and PIT-tag ID number were recorded for each individual. After tagging, the eels were placed in a recovery tank with aerated water before being moved to a fish holding tank. A few days after the tagging (between 1–6 days), the eels were transported by car to Ymsen (transport time: ca 4 h) and released at various sites along the shoreline of the lake.

During tagging event 4, in July 1999, $N=83$ yellow eels (mean size: 542 mm, Tab. 2) caught by a commercial fisher operating in Ymsen, were returned to the lake after PIT-tagging (note that the legal minimum catch-size was not applied since the fishing was performed for monitoring and research purposes). The tagging procedure and data collection were the same as for the tagging in 1998–2000 described above, with the difference that it was performed on-site (performed by HW and other staff at the Swedish Board of Fisheries). After tagging, the eels were kept in a fish corf for approximately two days to recover from the tagging, before being released into the lake.

During tagging event 5, in October 2012, $N=118$ silver eels (mean size: 859 mm, Tab. 2) caught by a commercial fisher operating in Ymsen, were returned to the lake after both Carlin- and PIT-tagging. The tagging procedure and data collection were performed in a similar way as described above for the PIT-tagging of yellow eels in July 1999, with the difference that the eel total mass was measured with ± 0.5 g precision, and these eels were tagged with both a Carlin- and a PIT-tag (performed by HW and JSt). The Carlin tag consisted of an 18 × 4 mm plastic plate (0.12 g, Inplastor AB, Motala, Sweden) with steel wires manually added to the plastic plate [by staff at Swedish University of Agricultural Sciences, Department of Aquatic Resources (SLU Aqua)]. The Carlin-tag was attached through the dorsal muscle below the dorsal fin with a double steel wire (while the eel was sedated, as described above for the PIT-tagging). Two injection needles were first disinfected by immersion in 95% ethanol, after which



Fig. 1. Photo of an adult female European eel, *Anguilla anguilla*, tagged with an external Carlin tag. Photo: Elin Myrenås.

they were pushed through the dorsal muscle approximately one centimetre below and two centimetre in front of the dorsal fin (Fig. 1). One steel wire thread was inserted through each injection needle, the needles were then removed, and the steel wires were wired together to lock the tag in place (Fig. 1). After tagging, the eels were kept in a recovery tank with freshwater until released in Ymsen the same day as they had been tagged.

2.3 Data collection and dissection, recaptures of Carlin and PIT-tagged eel

Data on recaptures of the PIT-tagged eels have been collected since 1999 (still ongoing in 2025). The only commercial fisher operating in Lake Ymsen continuously scans all captured eels in his catch for PIT tags. Eels were caught either in pound nets with 24 mm mesh-size or in an outlet trap with a 20 mm grid located in Ölebäcken (WGS84: 58.644841, 13.93384), which captures individuals migrating out of the lake. All scanned eels with a detected PIT-tag were handled in one of the following ways: a PIT-tagged eel from the 2012 tagging event was checked for the presence of a Carlin tag (since all PIT-tagged eels in 2012 also had a Carlin tag), and the presence or absence of the Carlin tag was noted. For those eels, no additional data was collected prior to their commercial sale. All other PIT-tagged eels (i.e., from any other PIT-tagging event in 1998–2000) were frozen in accordance with the commercial catch process, and sent for later dissection to SLU Aqua, except for 21 eels that were measured by the fisher as fresh, and the length and mass data was sent to SLU Aqua. Before dissection, the frozen eels were thawed. Length and mass (scale model: Mettler PC 400; Mettler Instrument AG, Küsnacht) were measured (mass was measured without removing coagulated mucus; Sundin et al., 2025). The abdomen was opened using scissors, from the anal opening

to the head. An ocular assessment of the gonads was made for sex determination (Tesch, 1977). All dissection data were stored in a database (Sötebasen), hosted by SLU Aqua, Institute of Freshwater Research. The dissections were performed by JP, JSt, supported by other staff at the Swedish Board of Fisheries/SLU Aqua.

2.4 Data collection, untagged fish (control)

To compare data from the recaptured tagged fish, we used data from untagged fish as a baseline control, which had been collected for monitoring purposes. Those eels had been sampled from the catch of the commercial fisher operating in Ymsen (using the same pound nets and outlet trap as described above) (2003: $n=85$; 2004: $n=30$; 2012: $n=11$; 2019: $n=123$; 2020: $n=129$; 2021: $n=130$, unknown catch year: $n=8$ [included in condition analysis]). From 2011 onward, all landed eels had a minimum size of 700 mm due to legal minimum catch-size restrictions. The sampled eels were frozen and sent to SLU Aqua for later dissection. The dissections were performed as described above, with the addition that the largest otoliths (sagittae, two per individual) were collected and saved for later age determination in a subset of the eels (according to methods described in ICES, 2009, with otoliths grinded, etched and stained) (untagged eels; aged: $n=440$, unaged: $n=76$). All dissection data were stored in the database Sötebasen.

2.5 Data analysis

All analyses were performed using R, version 4.2.1 (R Core Team, 2022) (by EM). Plots were constructed using the R-packages ggplot2 (Wickham et al., 2024) and cowplot (Wilke, 2024). Significance level was set to $p=0.05$. Language model software was used to generate and optimise

some of the R code drafts, to aid interpretation of R outputs, and language editing (OpenAI 2025; Google 2025). All AI-generated outputs were reviewed and edited by the authors.

For the Carlin and PIT-tagging studies, data validation had to be performed prior to analysis, since the eels in these studies were released a long time ago, and data collection on recaptures had been performed by several different staff members over time. Digital data in Excel spreadsheets was validated against original written protocols, ensuring that the data in the excel spreadsheets was correct and complete. PIT-tags that were stored together with samples (otoliths) from recaptured eels were scanned and checked against the digital data to ensure conformity. When validation and organisation of the data were completed (performed by EM, BJ, JSu, JSt, and PJ), some errors remained, such as certain recaptures missing release data. Data with unresolved errors were excluded from analyses.

Data on tagged and untagged (control $n = 516$) eel from Lake Ymsen were extracted from the database Sötebasen on August 20, 2025. Since most of the eels were measured as frozen and thawed, we corrected mass and length using a freeze-shrinking correction factor (Sundin et al., 2025) before further analyses.

The recapture rate was calculated as the proportion of individuals recaptured from the total number of released individuals per tagging event (Tab. 2). The mean time between the release event and recapture was calculated for PIT-tagged eels released as small yellow eels, for larger yellow eels, and for Carlin-tagged silver eels separately (Tab. 2). For Carlin tagged individuals (tagged at the silver eel stage), tag retention rate was calculated as the proportion of recaptured individuals retaining their tag from the total number of recaptured individuals.

Growth was evaluated by examining the relationship between length of the PIT-tagged eels and time (in years) elapsed since stocking (“years since release”) by fitting a LOESS (locally estimated scatterplot smoothing) curve for the two tagged groups separately: small and larger yellow eel, based on their size category at the time of stocking (Tab. 2). This non-parametric method fits multiple local regressions across the data range to produce a smooth trend without assuming a specific functional form. Each eel contributed with only two observations—length at release and length at recapture—and no repeated-measures data from the same individuals were available.

Specific growth rates (SGR) (Crane et al., 2020), expressed as annual change in percentage growth rate was calculated for the eels released as small yellow eels. The instantaneous growth rate constant (g) was first calculated from changes in mass (M) or length (L) between release date (t_1) and recapture date (t_2), for each eel, as:

$$g = (\ln(M_2) - \ln(M_1)) / (t_2 - t_1)$$

where M_1 and M_2 are the masses at release and recapture, respectively, and $(t_2 - t_1)$ is the time interval in months. An equivalent formula was used for length, replacing M with L . Then SGR was calculated by converting g from log scale to annual percentage change:

$$\text{SGR}(\% \text{ per year}) = (e^{(g^{12})} - 1) \cdot 100.$$

Table 3. Recaptures of Passive Integrated Transponder (PIT) -tagged European eels (*Anguilla anguilla*) released in Lake Ymsen in Sweden: release date of respective PIT-tagging event, year 1998–2000, tagging size of released eel (small yellow, or large yellow eel), number of PIT-tagged eels released, number of PIT-tagged eels recaptured, and recapture percentage.

Date released	N PIT-tag released	N recaptured	% recaptured
1998-10-28	190	0	0
1999-07-29	83	7	8
1999-09-15	1512	172	11
2000-09-14	1087	165	15

Three individuals tagged as small yellow eels were missing stocking- or catch data on mass and length and were excluded from these sgr-analyses.

To evaluate whether tagging influenced growth, we estimated the growth coefficient (k) and the asymptotic length (L_∞) parameters with Pütter-von Bertalanffy growth models (Beverton, 1954). We compared several candidate growth models to determine the model that best described growth differences between control and PIT-tagged small yellow eel. Growth models were fitted using nonlinear least squares (using the function `nls()` in R). Initial parameter values were obtained using the function `vbStarts()` from the FSA package (Ogle, 2013). The general model was defined as:

$$L(a; L_\infty, K, t_0) = L_\infty (1 - e^{-K(a-t_0)})$$

where $L(a)$ is the average length at age a , L_∞ is the asymptotic average length at which growth is zero, K is the growth rate coefficient (yr^{-1}), and t_0 , commonly interpreted as the hypothetical age at which predicted mean length equals zero. A set of candidate models was fitted, including a full model in which all parameters (L_∞ , K , and t_0) varied between groups and reduced models in which one or more parameters were shared. The model with the lowest Akaike’s Information Criterion (AIC) was selected as the best-supported model, using the function `AIC()`. For this, only aged untagged eels in the control group and eels tagged as small eels (age known due to tagging taking place in the first year after glass eel stage/import) were used, and no length at tagging was included in the data on which the models were fitted. Although batch (different stocking years of tagged eels) could not be included as a random effect in this model, preliminary model evaluations indicated that batch-related differences were limited to t_0 and did not affect estimates of growth rate (K) or asymptotic length (L_∞).

To evaluate differences in body condition between tagged and untagged eels, we fitted a linear model relating $\log_{10}$ - transformed mass to $\log_{10}$ - transformed length, treatment (PIT-tagged versus untagged) and their interaction ($\log_{10}$ length $\times$ treatment). Batch was not included as a random effect as it was undefined for wild individuals (control eels) and showed negligible variance in stocking year for restocked individuals (0.012% of the variance in $\log_{10}$ (mass) after accounting for $\log_{10}$ (length), $\eta = 0.00012$).




Fig. 2. Length differences between release and catch for PIT-tagged European eels (*Anguilla anguilla*) released in Lake Ymsen, Sweden, for A) released as small yellow eels (years 1998, 1999 and 2000), and B) released as larger yellow eels (year 1999). Total length for all eels when released is shown at year 0 (i.e., irrespective of actual release year) and recaptured as the number of years since release. For each individual that has been recaptured (PIT-tagged as small yellow $n = 334$, larger yellow eels $n = 7$), a line (grey) is shown between length (mm) at release and recapture. The solid black line represents a locally estimated scatterplot smoothing (LOESS) curve, which models the relationship between years since release and total length (mm), with the red shaded region representing the 95% confidence interval for the estimated curve. Panel C shows mean (± 1 SD) percentage increase in specific growth rates (SGR) for each year since release ($\Delta\text{Time} = \text{recapture date} - \text{release date}$), calculated based on mass (g year^{-1}) and length (mm year^{-1}), and the number of individuals (n) in each group. Only recaptured eels PIT-tagged as small yellow eels were included.

3 Results

3.1 Recaptured Carlin tagged silver eels

Out of the 118 individual silver eels that were tagged with both Carlin and PIT-tags and released in Ymsen in 2012, 76 (64%) have been recaptured (monitored until 2025). Records of the year of the catch were only available for 64 recaptures, but these were all recaptured within four years after being released [mean time until recaptured = 1 year (± 0.76)]. For the 76 recaptured eels, tag retention of the Carlin tag was 92% (restrictive number) or 97%, with data interpretation being the reason for the two numbers (due to issues regarding data quality on four recaptured individuals, with one source indicating that the tag was retained, while another source indicated that the tag was lost). None of the recaptured Carlin tagged eels had lost their PIT-tag. The calculations of tag retention rates disregard the unlikely possibility that both the Carlin and PIT-tags would have been lost (generally PIT-tags

have a high retention rate, Tab. 1), in which case the eel would be registered as untagged in the data.

3.2 Recaptured PIT-tagged yellow eels

Out of the 2872 PIT-tagged eels released in Ymsen in 1998–2000 (Tabs. 2 and 3), 344 (12%) have been recaptured (until 2025), with recapture rates varying among the different release batches (0–15%; Tab. 3). The eels released as small yellow eels were recaptured after 4–22 years (Fig. 2A) [$n = 337$, mean time between release and recapture = 12 years (± 4), with the longest possible duration being 27 years; 1998–2025]. The eels released as larger yellow eels (in summer, 1999) were recaptured after 3–15 years (Fig. 2B) [$n = 7$, mean time = 9 years (± 4)]. All recaptured PIT-tagged eels were females.

The relationship between mass or length and years spent in the lake after being released show large individual variation

Table 4. Summary table for the Pütter- von Bertalanffy growth model for PIT-tagged European eels and untagged control eels in Lake Ymsen. The table shows the model fit relative standard error (RSE), parameter estimates with their standard errors (SE), and the results of t -tests ($t(df)$, p -value) assessing parameter significance. “Group comparison” indicates the experimental groups, i.e., PIT- tagged as small yellow eels (TAG) vs.untagged control eels (C). Notes include details on parameter constraints or fitting remarks.

Group comparison	Model fit (RSE)	Parameter	Estimate $\pm$ SE	$t(df)$	p -value	Notes
TAG vs. C	63.66	K	0.23 ± 0.01	19.93 (770)	<0.001	Shared across groups
		L_{∞} (C)	956.79 ± 7.44 mm	128.62 (770)	<0.001	–
		L_{∞} (TAG)	924.68 ± 8.53 mm	108.34 (770)	<0.001	–
		t_0 (C)	3.91 ± 0.20 yr	19.62 (770)	<0.001	–
		t_0 (TAG)	1.85 ± 0.22 yr	8.45 (770)	<0.001	–




Fig. 3. Length at age data with Pütter-von Bertalanffy curves and 95% confidence intervals in shaded colours, for untagged control eels (black) ($n = 440$) and PIT-tagged eels released as small yellow eels (red) ($n = 335$). A minor horizontal jitter was added to the data points to enable better visualization of the data.




Fig. 4. Linear regressions of log-transformed mass (g) as dependent on log-transformed length (mm) (representing relative body condition) for Passive Integrated Transponder (PIT)-tagged European eels (*Anguilla anguilla*) released and later recaptured in Lake Ymsen (red data points and line), $n = 342$, and untagged, control eels captured in Lake Ymsen (black data points and line), $n = 516$. The shaded regions in grey and red represent the 95% confidence intervals for the regression lines (note that the 95% CI are very close to the regression line and therefore difficult to see in the figure).

Table 5. Summary table for the linear model of log-transformed mass (M) as dependent on log-transformed length (TL) (i.e., relative body condition) for Passive Integrated Transponder (PIT)-tagged European eels (*Anguilla anguilla*) ($n = 342$) released and recaptured in Lake Ymsen. The reference group (intercept) was the untagged eels, the control group (from Lake Ymsen) ($n = 516$).

Response variable	Coefficient	Estimate (β)	SE	T	p
$\log_{10}(M)$	Intercept	−6.49	0.05	−130.73	<0.001
	$\log_{10}(TL)$	3.27	0.02	191.41	<0.001
	Treatment [PIT]	0.04	0.11	0.39	0.70
	$\log_{10}(TL) \times \text{treatment [PIT]}$	−0.01	0.04	−0.37	0.71

both for eels released as small yellow eels (Figs. 2A and 2C) and as larger yellow eels (Fig. 2B).

The growth of eels was analyzed using Pütter-von Bertalanffy growth models, with parameters estimated separately for different tagging groups (Tab. 4).

The growth of eels tagged with PIT-tags as small yellow eels and untagged control eels was modelled with separate asymptotic lengths (L_{∞}) and theoretical ages at zero length (t_0) for the two groups, and a shared growth rate parameter (K) across groups, as this model got the smallest AIC. The next-best model, which also allowed the growth parameter (K) to vary between groups, increased model complexity ($\Delta\text{AIC} \approx 2$). The estimated asymptotic length (L_{∞}) was higher (~ 32 mm) for the untagged control group compared to the tagged group (Tab. 4). The theoretical age at zero length (t_0) also differed between the two groups; the tagged eels had a lower estimated t_0 (Tab. 4). These results suggest that both groups grew at the same rate but differed in their initial size and asymptotic length (Fig. 3A).

There were no significant differences in either intercept or slope between PIT-tagged eels and untagged control eels in mass corrected for length, i.e. relative body condition ($F_{3,854} = 1.52 \cdot 10^4$, $p < 0.001$) (Fig. 4, Tab. 5).

4 Discussion

In this study, we investigated tag retention of Carlin tags in silver eels, and any potential effects on growth and condition of PIT-tagged yellow eels by utilising data from several long-term field studies conducted in a natural lake in Sweden. We found that the tag retention rate of the recaptured Carlin tagged eels was 92%, with six out of the 76 recaptured eels having lost their Carlin tag. The retention rate could be as high as 97%, given uncertainties between different data sources regarding four recaptured individuals. Our results are in line with a laboratory study performed by Vøllestad (1988), where he could see a retention rate of Carlin tags at 90%. In general, certain behaviours of fish, such as passing through crevices and burrowing, can lead to loss of external tags (Jepsen et al., 2015). Since the eel is a bottom-dwelling fish that inhabits hollows and burrows, sometimes several eels together, and can spend a long time within the sediment during winter dormancy (Nyman, 1972; Tesch, 1977; Westerberg and Sjöberg, 2014), this could be considered as one potential factor underlying the 8% Carlin-tag loss. The study by Vøllestad (1988) only monitored tagged eels during seven months in a laboratory environment, hence possibly not covering tag losses due to such types of behaviour. With a possible recapture period of

more than twenty years in our study, any seasonal variation in tag loss should be sufficiently covered. The comparable retention rates between the studies suggest that burrowing is not the primary cause of tag loss. If the eel would get entangled and lose the tag in the fishing gear, there would likely be a wound noted during the dissection. But there are other specific behaviours of the eel, like the ability to tie their body into a knot, rolling around, or reaching and biting at the tag, that could mean that the eel might not be an ideal species for external tags (Økland et al., 2013; Jepsen et al., 2015). The retention rate has been found to be as low as 0–9% of larger external telemetry tags on eels (Cottrill et al., 2006; Økland et al., 2013), and these types of behaviour might be the reason for tag loss of Carlin tags as well. Considering the very low retention rate of larger tags, the tag retention of Carlin tags at 92%, as found in this study, could be considered high. In precaution, when accounting for retention rate in statistical models and analyses using recaptures of Carlin tagged eel, the use of the lower retention rate found in this study (92%) is recommended, with additional research encouraged to strengthen confidence in these estimates. Particularly, we encourage more studies across different habitats, since eels can inhabit both lakes and coastal environments as well as migrate through more turbulent rivers between these habitats (Tzeng et al., 1997; Teichert et al., 2023), potentially exposing the tag to additional influencing factors beyond those considered in this study.

All recaptured Carlin tagged eels with known recapture date were recaptured within four years after being released, despite continued monitoring for 27 years (continuing in 2025). A possible explanation could be that the combination of the fishing gear and outlet trap have not captured 100% of the migrating eel, or that some of the remaining tagged individuals died. Vøllestad (1988) found a mortality rate of 13% in the laboratory study monitoring Carlin-tagged eel, which would correspond to 15 individuals in our study, much lower than the 43 individuals that were not recaptured (corresponding to a mortality rate of 36%). There are several reasons why tagging might lead to mortality: direct (e.g. infections) or indirect effects (e.g. increased predation due to a visible external tag) of the Carlin tag, and these might be higher in a natural environment than in a laboratory. Tag-related mortality was, however, not possible to evaluate in this study, and similarly, natural mortality of untagged eels cannot be evaluated here. Tags should ideally not influence the survival, growth or behaviour of the tagged individuals and should last as long as possible to fulfil their purpose, which is true regardless of tag type. While Carlin tags currently seem to be less commonly used than they have been in the past, it is still an important tool

to estimate fishing mortality for eel stock assessment purposes in Sweden (van Gemert et al., 2024). In the long term, the availability of robust stock assessments is crucial for implementing effective management actions that support population recovery and allow for sustainable use of the resource. Additionally, Carlin-style tags are currently used in other species of fish in scientific studies, for example on lake trout (*Salvelinus namaycush*) (Guzzo et al. 2025) and Atlantic salmon (*Salmo salar*) (Sheehan et al. 2024). Hence, to ensure correct interpretation of tagging data, future work should investigate the mortality rate caused by Carlin tagging, specifically to separate different possible adverse effects, like infections and increased predation pressure, from natural mortality.

Our results regarding the effects of PIT-tags add to the collection of studies that do not find any major impacts of PIT-tags on growth or condition in eels. In general, the PIT-tagged eels were found to have a similar growth rate as untagged eels. Although the von Bertalanffy growth model indicated differences in asymptotic length (L_{∞}) between control and those tagged as small yellow eels, these differences should be interpreted cautiously. The estimated difference in asymptotic length (L_{∞}) of 32 mm (control: 957 mm, tagged: 925 mm) is modest when viewed in the context of the natural size variability commonly reported for silver eels in Sweden (Svedäng et al., 1996; Figure C14 in van Gemert et al. 2024). Importantly, the growth rate (K) did not differ between tagged and untagged eels, indicating that PIT-tagging did not impair somatic growth. The conclusion that PIT-tags do not have any clear effects on eels was further strengthened since there was no significant difference in relative condition (length-mass relationship) between tagged and untagged eels in our study. There were, however, no recaptures from the first stocking year of PIT-tagged eels in Ymsen in 1998, and the reason for that is unknown. The number of released PIT-tagged eels that year was rather low, $n = 190$, in particular compared to the other two years when small yellow eels were PIT-tagged and restocked ($n = 1512$ and 1087 , respectively, in 1999 and 2000). However, there were 7 recaptured eels from the even smaller summer batch of larger tagged yellow eels in 1999, $n = 83$. These tagged yellow eels were large enough to get caught in the fishing gears relatively soon after being released, while the tagged small yellow eels from the batch in 1998 had to grow for a longer period before being large enough to possibly be caught in the gears. Importantly, if any of those eels were male, they would have migrated from the lake at a size too small to be caught in the fishing gear, since males are much smaller than females (Vøllestad and Jonsson, 1986). An important aspect to consider in tagging studies that rely on recapture reports by fishers is the uncertainty associated with the fishers' recognising the tag and/or the willingness to report having captured a tagged fish (Pollock et al., 2001). There is no reason to believe that the cooperation with the fisher in Lake Ymsen is anything but good, since the communication over the years has worked well and since small-scale fisheries in general may feel a higher responsibility for their work and resource than large-scale fisheries (Berkes et al., 2001). The lack of recaptures from the stocking in 1998 is more likely explained by some of those eels being males and not being recaptured, and/or there could have been a

methodological problem during tagging or health issues in the supplied eel, even though no such reason is known.

An important and unusual aspect of this study is that the fisher has reported catches from both fishing gear and the outlet trap for more than 27 years (still ongoing in 2025). Only a few recaptures of PIT-tagged eels have been made in recent years, the most recent in 2022. Most of the tagged eels are expected to have matured into silver eels during this period and should have a natural urge to leave the system (hence being caught in the fishing gear or in the outlet trap), although eels in the northern hemisphere can reach a considerable age (30 years or more, Durif et al., 2020). Consequently, this long-term field study likely covers the majority of the yellow eel growth phase of the released PIT-tagged eels (from restocked small yellow eel to mature silver eel), under the rare conditions of a well-controlled natural lake system. The recapture rate of PIT-tagged eels (on average 12%) was lower than for Carlin-tagged eels (64% recapture rate). This is, however, not surprising since the eels tagged with Carlin tags were large and most likely old eels. Compared to young individuals, older fish can have a substantially lower natural mortality (Bevacqua et al., 2011) and a higher natural urge to leave the system, leading them to the outlet trap or fishing gear (i.e. being recaptured). From 2011 onward, catches from the commercial fisheries had a size limit of 700 mm, which could have biased recapture rates. However, given the long study period, allowing eels to grow and age, increasing the possibility of being caught in the fishing gear or the outlet trap, this effect is likely minimal. Compared to other long-term field studies with PIT-tagged young eels, the recapture rate (12%) lies at the same level as for both tagged and untagged eels. One example is a study performed in France where they found a 15% recapture rate of PIT-tagged eels after 11 years in a natural system (Mazel et al., 2013). Another example is a study of untagged restocked eels in a Swedish lake (the lake had no eels prior to the restocking event), where 11% had been recaptured in the outlet trap after 14 years (Wickström et al., 1996). Overall, our results suggest no clear negative effect of PIT-tagging on the survival of yellow eel in natural lake systems, despite survival not being directly assessed in this study.

5 Conclusion

Based on the relatively high retention rate of Carlin tags and the lack of negative effects on condition or growth rates of PIT-tags, we conclude that these are suitable methods for tagging European eel. For the Carlin tags, the results only regard silver eels, and these kinds of tags might not be as suitable for yellow eels, which have less thick skin. In addition, there are other aspects of Carlin tagging that are not covered in this study that also need to be considered in tagging studies, such as tagging mortality. While the results from this study regarding Carlin tags can significantly contribute to improving stock assessments and mortality estimates based on mark-recapture studies using Carlin tags, the results regarding PIT-tags can further improve meta-analyses and welfare legislation. This, in turn, may facilitate the implementation of appropriate management strategies and conservation endeavours for the critically endangered European eel that promote population recovery and long-term sustainable fisheries in the future.

Acknowledgments

We thank the local fishers, the family Grönlind, who helped with the field PIT- and Carlin tagging studies conducted in Lake Ymsen, by monitoring all eels caught in Lake Ymsen for the presence of PIT- and/or Carlin tags since 1999. We thank the staff of the former Swedish Board of Fisheries, and the present Swedish University of Agricultural Sciences, Department of Aquatic Resources (SLU Aqua), Institute of Freshwater Research, for their assistance with tagging, animal care, data handling, dissections, and long-term data collection. Their efforts have contributed to the important and substantial database “Sötebasen”, which made this study possible. Thanks to Scandinavian Silver Eel (SSE) for assistance with obtaining the juvenile eels and providing work premises during the tagging procedure. We also thank the reviewers and editors for their constructive comments and suggestions, which helped improve the quality and clarity of this manuscript.

Funding

Given the long-time span of the data used in the study, the data have been collected as part of several different monitoring programs, with the most recent being the EU data collection framework in the fisheries and aquaculture sector (DCF, Regulation 2017/1004), partially funded by the EU.

Conflicts of interest

The authors declare no competing interests. The authors have no relevant financial or non-financial interests to disclose.

Data availability statement

The data and script for this study are archived in the figshare repository: <https://doi.org/10.6084/m9.figshare.30731642>, and <https://doi.org/10.6084/m9.figshare.30731711> following best practices (Roche et al., 2015) and were available to editors and reviewers upon initial submission.

Author contribution statement

Author contribution roles are listed here according to the Contributor Role Taxonomy (CRediT) guidelines (<https://credit.niso.org>): Conceptualization: JSu, HW, JN, EM; Investigation: EM, JN, JP, HW, JSt, JSu; Data curation: JSt, EM, JN, BJ, JP, PJ; Formal analysis: EM, JN; Funding acquisition: HW, JSu; Project administration: JSu, HW; Validation: EM, BJ, JN; Visualization: JN, EM; Writing - original draft: EM, JSu; Writing - review & editing: all authors.

Ethical statement

The PIT- and Carlin tagging performed in 1998–2000 and 2012 were performed in accordance with the animal ethical legislation valid at that time. All recaptured eels used in this study originated from a commercial fishery that complied with fisheries regulations.

References

Acou A, Boury P, Laffaille P, Crivelli AJ, Feunteun E. 2005. Towards a standardized characterization of the potentially migrating silver

European eel (*Anguilla anguilla* L.). *Archiv für Hydrobiologie*, 164: 237–255.

Acou A, Feunteun E, Laffaille P, Legault A. 2000. Catadromous migration dynamics of European eel (*Anguilla anguilla*, L.) in a dammed catchment. *Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen*, 27: 3117–3120.

Atkins CG. The Atlantic salmon (*Salmo salar*). A—Report on the collection and distribution of Penobscot salmon in 1873–’74 and 1874–’75, United States Commission of Fish and Fisheries, Washington, 1876, pp 485–540.

Berkes F, Mahon R, McConney P, Pollnac R, Pomeroy R. Managing Small-Scale Fisheries, Alternative Directions and Methods, International Development Research Centre, Ottawa, Canada, 2001. ISBN 0-88936-943-7

Bevacqua D, Melia P, De Leo GA, Gatto M. 2011. Intra-specific scaling of natural mortality in fish: the paradigmatic case of the European eel. *Oecologia* 165: 333–339.

Beverton RJH. Notes on the Use of Theoretical Models in the Study of the Dynamics of Exploited Fish Populations, Miscellaneous Contributions Number 2. U.S. Fishery Laboratory, Beaufort, NC, 1954.

Bridger CJ, Booth RK. 2003. The effects of biotelemetry transmitter presence and attachment procedures on fish physiology and behavior. *Rev Fish Sci* 11: 13–34.

Cahill CL, Mogensen S, Wilson KL, Cantin A, Sinnatamby RN, Paul AJ, Christensen P, Reilly JR, Winkel L, Farineau A, Post JR. 2018. Multiple challenges confront a high-effort inland recreational fishery in decline. *Can J Fish Aquat Sci* 75: 1357–1368.

Carlin B. 1955. Tagging of salmon smolts in the River Lagan. *Rep Inst Freshw Res Drottningholm* 36: 57–74.

Carrier JC. 1985. Nurse sharks of Big Pine Key: comparative success of three types of external tags. *Florida Sci* 146–154.

Center NF. 1990. Equipment, methods, and an automated data-entry station for PIT tagging. *Am Fish Soc Symp* 7: 335–340.

Cottrill RA, Økland F, Aarestrup K, Jepsen N, Koed A, Hunter KJ, Butterworth KG, McKinley RS. 2006. Evaluation of three telemetry transmitter attachment methods for female silver-phase American eels (*Anguilla rostrata* Lesueur). *J Great Lakes Res* 32: 502–11.

Cowen L, Schwarz CJ. 2005. Capture–recapture studies using radio telemetry with premature radio-tag failure. *Biometrics* 61: 657–664.

Crane DP, Ogle DH, Shoup DE. 2020. Use and misuse of a common growth metric: guidance for appropriately calculating and reporting specific growth rate. *Rev Aquac* 12: 1542–1547.

Cucherousset J, Roussel JM, Keeler R, Cunjak RA, Stump R. 2005. The use of two new portable 12-mm PIT tag detectors to track small fish in shallow streams. *North Am J Fish Manag* 25: 270–274.

Dainys J, Gorfine H, Šidagytė E, Jakubavičiūtė E, Kirka M, Putys Ž, Ložys L. 2017. Do young on-grown eels, *Anguilla anguilla* (Linnaeus, 1758), outperform glass eels after transition to a natural prey diet? *J Appl Ichthyol* 33: 361–365.

Dekker W, Sjöberg NB. 2013. Assessment of the fishing impact on the silver eel stock in the Baltic using survival analysis. *Can J Fish Aquat Sci* 70: 1673–1684.

Durif CM, Diserud OH, Sandlund OT, Thorstad EB, Poole R, Bergesen K, Vøllestad LA. 2020. Age of European silver eels during a period of declining abundance in Norway. *Ecol Evol* 10: 4801–4815.

Eames MJ, Hino MK. 1983. An evaluation of four tags suitable for marking juvenile chinook salmon. *Trans Am Fish Soc* 112: 464–468.

- Elliott DG, Pascho RJ. 2001. Evidence that coded-wire-tagging procedures can enhance transmission of *Renibacterium salmoninarum* in Chinook salmon. *J Aquat Anim Health* 13: 181–193.
- Guzzo MM, Blanchfield PJ, MacLeod HA, Rennie MD. 2025. Declining adult body size and contracted life history in an unexploited boreal lake trout (*Salvelinus namaycush*) population during a period of warming. *Can J Fish Aquat Sci* 82: 1–15.
- Hays GC, Bailey H, Bograd SJ, Bowen WD, Campagna C, Carmichael RH, Casale P, Chiaradia A, Costa DP, Cuevas E, de Bruyn PJN, Dias MP, Duarte CM, Dunn DC, Dutton PH, Esteban N, Friedlaender A, Goetz KT, Godley BJ, Sequeira AMM. 2019. Translating marine animal tracking data into conservation policy and management. *Trends Ecol Evol* 34: 459–473.
- Herrera M, Moreno-Valcárcel R, De Miguel Rubio R, Fernández-Delgado C. 2019. From transient to sedentary? Changes in the home range size and environmental patterns of movements of European eels (*Anguilla anguilla*) in a Mediterranean River. *Fishes* 4: 43.
- Hirt-Chabbert JA, Young OA. 2012. Effects of surgically implanted PIT tags on growth, survival and tag retention of yellow shortfin eels *Anguilla australis* under laboratory conditions. *J Fish Biol* 81: 314–319.
- Höhne L, Freese M, Pohlmann JD, Diekmann M, Fladung E, Huisman JB, Hanel R, Marohn L. 2023. Overestimating management progress—modelled vs. monitored silver eel escapement in a North Sea draining river. *ICES J Mar Sci* 80: 1936–1948.
- Holmgren K, Mosegaard H. 1996. Implications of individual growth status on the future sex of the European eel. *J Fish Biol* 49: 910–925.
- Huusko A, Mäki-Petäys A, Stickler M, Mykrä H. 2011. Fish can shrink under harsh living conditions. *Funct Ecol* 25: 628–633.
- Huusko R, Huusko A, Mäki-Petäys A, Orell P, Erkinaro J. 2016. Effects of tagging on migration behaviour, survival and growth of hatchery-reared Atlantic salmon smolts. *Fish Manag Ecol* 23: 367–375.
- ICES. Workshop on Age Reading of European and American Eel (WKAREA), 20–24 April 2009, Bordeaux, France, 2009, ICES CM 2009/ACOM: 48.
- ICES. Report of the Joint EIFAAC/ICES/GFCM Working Group on Eel (WGEEEL). 24 November–2 December 2015, Antalya, Turkey, 2015, ICES CM 2015/ACOM:18.
- Jacobson B, Wickström H, Strömquist J, Persson J. Ålmärkning-sprojekt i sjön Ymsen; sammanställning av data från 1998–2022. Aqua notes 2024:3, Institutionen för akvatiska resurser, Uppsala, 2024.
- Jellyman DJ, Crow SK. 2016. Population size, growth and movements of *Anguilla australis* in a small lake. *J Fish Biol* 88: 2157–2174.
- Jepsen N, Richter L, Pedersen MI, Deng Z. 2022. Survival, growth and tag retention of juvenile European eel (*Anguilla anguilla* L.) with implanted 12 mm passive integrated transponder tags and acoustic tags. *J Fish Biol* 101: 1375–1380.
- Jepsen N, Schreck C, Clements S, Thorstad EB. 2005. A brief discussion on the 2% tag/bodymass rule of thumb. *Aquatic Telemetry: Adv Appl*, 255–259.
- Jepsen N, Thorstad EB, Havn T, Lucas MC. 2015. The use of external electronic tags on fish: an evaluation of tag retention and tagging effects. *Anim Biotelemetry* 3: 1–23.
- Kallemeyn LW. 1989. Loss of Carlin tags from walleyes. *North Am J Fish Manag* 9: 112–115.
- Kerstetter DW, Polovina J, Graves JE. 2004. Evidence of shark predation and scavenging on fishes equipped with pop-up satellite archival tags. *Fish Bull* 102: 750.
- Kushnir D, Degani G. 1991. Growth performance of European eel (*Anguilla anguilla*) under controlled photoperiod and shelter availability. *Aquac Eng* 10: 219–226.
- Laffaille P, Acou A, Guillouët J. 2005. The yellow European eel (*Anguilla anguilla* L.) may adopt a sedentary lifestyle in inland freshwaters. *Ecol Freshw Fish* 14: 191–196.
- Larsen MH, Thorn AN, Skov C, Aarestrup K. 2013. Effects of passive integrated transponder tags on survival and growth of juvenile Atlantic salmon *Salmo salar*. *Anim Biotelemetry* 1: 1–8.
- Latour RJ. Tagging methods and associated data analysis, in: J.A. Musick, R. Bonfil (Eds.), *Management Techniques for Elasmobranch Fisheries*, FAO, Rome, 2005, pp. 45–61.
- Lucas MC, Baras E. 2008. Methods for studying spatial behaviour of freshwater fishes in the natural environment. *Fish Fish* 1: 283–316.
- Mazel V, Charrier F, Legault A, Laffaille P. 2013. Long-term effects of passive integrated transponder tagging (PIT tags) on the growth of the yellow European eel (*Anguilla anguilla* (Linnaeus, 1758)). *J Appl Ichthyol* 29: 906–908.
- Miljödata-MVM 2026. Swedish University of Agricultural Sciences (SLU). National Data Host Lakes and Watercourses, and National Data Host Agricultural Land, <https://miljodata.slu.se/mvm/> [2026-02-05].
- Miller MJ, Westerberg H, Sparholt H, Wysujack K, Sørensen SR, Marohn L, Jacobsen MW, Freese M, Ayala DJ, Pohlmann J-D. 2019. Spawning by the European eel across 2000 km of the Sargasso Sea. *Biol Lett* 15: 20180835.
- Morrison WE, Secor DH. 2003. Demographic attributes of yellow-phase American eels (*Anguilla rostrata*) in the Hudson River estuary. *Can J Fish Aquat Sci* 60: 1487–1501.
- Nakagawa S, Ivimey-Cook ER, Grainger MJ, O’Dea RE, Burke S, Drobnjak SM, ... Lagisz M. 2023. Method reporting with initials for transparency (MeRIT) promotes more granularity and accountability for author contributions. *Nat Commun* 14: 1788.
- Nielsen J. 1988. A review on marking and tagging methods applied to eel (*Anguilla anguilla* (L.)). *FAO EIFAC Occasional Papers*, 21: 24.
- Nordqvist H. 1928. Försök rörande import och inplantering i svenska sjöar av engelsk glasäl. *Skrifter utgivna av Södra Sveriges Fiskeriförening*, 1: 60–68.
- Nyman L. 1972. Some effects of temperature on eel (*Anguilla*) behaviour. *Rep Inst Freshw Res Drottningholm* 52: 90–102.
- Nyqvist D, Schiavon A, Candiotto A, Tarena F, Comoglio C. 2023. Survival and swimming performance in small-sized South European Cypriniformes tagged with passive integrated transponders. *J Ecohydraulics* 9: 248–258.
- Nyström A, Trybom F. 1902. Uppgångsrännor för ålyngel vid Trollhättan. *Svensk Fiskeritidskrift*, 11: 1–9.
- Nzau Matondo B, Ovidio M. 2018. Decreased stock entering the Belgian Meuse is associated with the loss of colonisation behaviour in yellow-phase European eels. *Aquat Living Resour* 31: 7.
- Nzau Matondo B, Benitez JP, Dierckx A, Renardy S, Rollin X, Colson D, Baltus L, Romain VRM, Ovidio M. 2021. What are the best upland river characteristics for glass eel restocking practice? *Sci Total Environ* 784: 147042.
- Ogle D. fishR vignette – von Bertalanffy Growth Models [R package vignette], 2013, Retrieved from <http://derekeogle.com/fishR/examples/oldFishRVignettes/VonBertalanffy.pdf>
- Økland F, Thorstad EB, Westerberg H, Aarestrup K, Metcalfe JD. 2013. Development and testing of attachment methods for pop-up

- satellite archival transmitters in European eel. *Anim Biotelemetry* 1: 1–13.
- Olivereau M, Olivereau JM. 1997. Long-term starvation in the European eel: general effects and responses of pituitary growth hormone-(GH) and somatolactin-(SL) secreting cells. *Fish Physiol Biochem* 17: 261–269.
- Pedersen MI, Dieperink C. 2000. Fishing mortality on silver eels, *Anguilla anguilla* (L.), in Denmark. *Dana* 12: 77–82.
- Pike C, Crook V, Gollock M. *Anguilla anguilla*. The IUCN Red List of Threatened Species 2020: e.T60344A152845178, 2020, <https://doi.org/10.2305/IUCN.UK.2020-2.RLTS.T60344A152845178.en>. Accessed on 20 February 2024.
- Piper AT, Wright RM, Walker AM, Kemp PS. 2013. Escapement, route choice, barrier passage and entrainment of seaward migrating European eel, *Anguilla anguilla*, within a highly regulated lowland river. *Ecol Eng* 57: 88–96.
- Pollock KH, Hoenig JM, Hearn WS, Calingaert B. 2001. Tag reporting rate estimation: 1. An evaluation of the high-reward tagging method. *North Am J Fish Manag* 21: 521–532.
- Prentice EF, Park DL. 1984. A study to determine the biological feasibility of a new fish tagging system. *Annual Report of Research* (1983-84), Project 83-19. National Marine Fisheries Service, Seattle, WA, 1983.
- Prentice EF, Flagg TA, McCutcheon CA, Brastow DF, Cross DC. 1990. Equipment, methods and automated data-entry station for PIT tagging. *Am Fish Soc Symp* 7: 335–340.
- R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, 2022. URL <https://www.R-project.org/>
- Riley WD, Walker AM, Bendall B, Ives MJ. 2011. Movements of the European eel (*Anguilla anguilla*) in a chalk stream. *Ecol Freshw Fish* 20: 628–635.
- Roche DG, Kruuk LEB, Lanfear R, Binning SA. 2015. Public data archiving in ecology and evolution: how well are we doing? *PLoS Biol* 13: e1002295.
- Saville A, Morrison JA. 1985. Evaluation of tagging and detection using microwave tags in herring and the potential of this technique in herring stock assessment. *ICES CM* 1985/H:26.
- Schmidt J. 1912. The reproduction and spawning-places of the freshwater eel (*Anguilla Vulgaris*). *Nature* 89: 633–636.
- Schmucker AK, Johnson NS, Galbraith HS, Li W. 2017. An evaluation of silver-stage American eel conspecific chemical cueing during outmigration. *Environ Biol Fishes* 100: 851–864.
- Semple SL, Mulder IM, Rodriguez-Ramos T, Power M, Dixon B. 2018. Long-term implantation of acoustic transmitters induces chronic inflammatory cytokine expression in adult rainbow trout (*Oncorhynchus mykiss*). *Vet Immunol Immunopathol* 205: 1–9.
- Sheehan TF, Gargan P, Kerr B, Nevoux M, Ravn M, Nygaard R, Bradbury IR, Robertson MJ, Ó Maoiléidigh N. 2024. The international sampling program: continent of origin and biological characteristics of Atlantic salmon (*Salmo salar*) collected at West Greenland in 2020 and 2021.
- Simon J, Dörner H. 2014. Survival and growth of European eels stocked as glass-and farm-sourced eels in five lakes in the first years after stocking. *Ecol Freshw Fish* 23: 40–48.
- Sjöberg NB, Petersson E, Wickström H, Hansson S. 2009. Effects of the swimbladder parasite *Anguillicola crassus* on the migration of European silver eels *Anguilla anguilla* in the Baltic Sea. *J Fish Biol* 74: 2158–2170.
- SLU Aqua. Database Sötebasen, 2026. [https://aqua.slu.se/al/\[2026-02-11\]](https://aqua.slu.se/al/[2026-02-11]).
- Sundin J, Persson J, Skoglund S. 2025. A correcting factor for the reduction of body length and mass of European eel after ethanol preservation and after freezing. *Aquac Fish Fisheries* 5: e70094.
- Svedäng H, Neuman E, Wickström H. 1996. Maturation patterns in female European eel: age and size at the silver eel stage. *J Fish Biol* 48: 342–351.
- Teichert N, Bourillon B, Suzuki K, et al. 2023. Biogeographical snapshot of life-history traits of European silver eels: insights from otolith microchemistry. *Aquat Sci* 85. <https://doi.org/10.1007/s00027-023-00940-4>.
- Tesch FW. 1989. Changes in swimming depth and direction of silver eels (*Anguilla anguilla* L.) from the continental shelf to the deep sea. *Aquat Living Resour* 2: 9–20.
- Tesch F-W. The Eel: Biology and Management of Anguillid Eels, Chapman and Hall, London, 1977.
- Thorstad EB, Rikardsen AH, Alp A, Økland F. 2013. The use of electronic tags in fish research—an overview of fish telemetry methods. *Turk J Fish Aquat Sci* 13: 881–896.
- Tzeng WN, Severin KP, Wickström H. 1997. Use of otolith microchemistry to investigate the environmental history of European eel *Anguilla anguilla*. *Mar Ecol Prog Ser* 149: 73–81.
- Vallin S. 1929. Sjöen Ymsen i Skaraborgs län. Meddelanden från Kungl. Lantbruksstyrelsen 277(6:1929): 1–43.
- van Gemert R, Holliland P, Karlsson K, Sjöberg N, Säterberg T. Assessment of the Eel Stock in Sweden, Spring 2024; Fifth Post-Evaluation of the Swedish Eel management. Aqua reports 2024:5, Swedish University of Agricultural Sciences (SLU), Uppsala, 2024, <https://doi.org/10.54612/a.4iseib7eup>
- Vandergoot CS, Brenden TO. 2014. Spatially varying population demographics and fishery characteristics of Lake Erie walleyes inferred from a long-term tag recovery study. *Trans Am Fish Soc* 143: 188–204.
- Verdon R, Desrochers D. 2003. Upstream migratory movements of American eel *Anguilla rostrata* between the Beauharnois and Moses-Saunders power dams on the St. Lawrence River. *Am Fish Soc Sympos* 33: 139–151.
- Verdon R, Desrochers D, Dumont P. 2003. Recruitment of American eels in the Richelieu river and Lake Champlain: Provision of upstream passage as a regional-scale solution to a large-scale problem. *Am Fish Soc Sympos* 33: 125–138.
- Vøllestad LA, Jonsson B. 1986. Life-history characteristics of the European Eel *Anguilla anguilla* in the Imsa River, Norway. *Trans Am Fish Soc* 115: 864–871.
- Vøllestad LA. 1988. Tagging experiments with yellow eel, *Anguilla anguilla* (L.), in brackish water in Norway. *Sarsia* 73: 157–161.
- Vollset KW, Lennox RJ, Thorstad EB, Auerm S, Bär K, Larsen MH, Mahlum S, Näslund J, Stryhn H, Dohoo I. 2020. Systematic review and meta-analysis of PIT tagging effects on mortality and growth of juvenile salmonids. *Rev Fish Biol Fish* 30: 553–568.
- Von Bertalanffy L. 1938. A quantitative theory of organic growth (inquiries on growth laws. II). *Human Biol* 10: 181–213.
- Westerberg H, Sjöberg N. 2014. Overwintering dormancy behaviour of the European eel (*Anguilla anguilla* L.) in a large lake. *Ecol Freshw Fish* 24: 532–543.

- Westerberg H, Sjöberg N, Lagenfelt I, Aarestrup K, Righton D. 2014. Behaviour of stocked and naturally recruited European eels during migration. *Mar Ecol Prog Ser* 496: 145–157.
- Westin L. 1990. Orientation mechanisms in migrating European silver eel (*Anguilla anguilla*): temperature and olfaction. *Marine Biol* 106: 175–179.
- Wickham H, Chang W, Pedersen TL, Takahashi K, Wilke C, Woo K, Yutani H, Dunnington D, van den Brand T. ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics, CRAN, 2024. <https://doi.org/10.32614/CRAN.package.ggplot2>
- Wickström H, Sjöberg NB. 2014. Traceability of stocked eels—the Swedish approach. *Ecol Freshw Fish* 23: 33–39.
- Wickström H, Westin L, Clevestam P. 1996. The biological and economic yield from a long-term eel-stocking experiment. *Ecol Freshw Fish* 5: 140–147.
- Wilke CO. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2', CRAN, 2024. <https://doi.org/10.32614/CRAN.package.cowplot>
- Wood SN. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J R Stat Soc: Ser B (Stat Methodol)* 73: 3–36.
- Wright GV, Wright RM, Kemp PS. 2015. Impact of tide gates on the migration of adult European eels, *Anguilla anguilla*. *Estuaries Coasts* 38: 2031–2043.
- Wright RM, Piper AT, Aarestrup K, Azevedo JM, Cowan G, Don A, Gollock M, Ramallo SR, Velterop R, Walker A, Westerberg H, Righton D. 2022. First direct evidence of adult European eels migrating to their breeding place in the Sargasso Sea. *Sci Rep* 12: 15362.
- Závorka L, Aymes J-C, Guiheneuf A, Mercier O, Huger F, Bejean M, Glisne S, Huchet E, Tomanova S. 2024. Effect of 12 mm PIT tags on the survival, growth, and ecologically significant behaviours of juvenile critically endangered endemic *Zingel asper*. *Knowl Manag Aquat Ecosyst* 425: 20.
- Zimmerman JL, Welsh SA. 2008. PIT tag retention in small (205-370 mm) American eels, *Anguilla rostrata*. *Proc West Virginia Acad Sci* 79: 1–8.
- Zimmerman JL. 2008. Rates of swim bladder parasite infection and PIT tag retention in upstream migrant American eels of the Upper Potomac River drainage [master's thesis]. Morgantown (WV): West Virginia University

Cite this article as: Myrenås E, Näslund J, Jacobson P, Jacobson B, Persson J, Wickström H, Strömquist J, Sundin J. 2026. High retention rate of Carlin tags and no effects on growth and condition of PIT tagging - a long-term field study on European eel. *Aquat. Living Resour.* 39: 20. <https://doi.org/10.1051/alr/2026015>